

Evaluation of Clinical Profile and Effect of Cervical Spinal Cord Injuries with or Without Bony Injuries

Sushil Acharya¹, Vinod Sharma¹, Sushil Sharma², Jitendra Acharya³

¹Associate Professor, Department of Neurosurgery, SMS Medical College, Jaipur, Rajasthan, India. ²Intern, S. P. Medical College, Bikaner, Rajasthan, India.

³Senior Demonstrator, Department of Dentistry, S.P. Medical College, Bikaner Rajasthan, India.

Abstract

Background: This study was design to study the clinical profile and outcome in patients of cervical spinal cord injury with or without bony injury. **Subjects and Methods:** All patients of cervical spinal cord injury with or without evidence of bony injury on X-ray were included in this study. In addition to X-ray finding, patient age, sex and MRI cervical spine were also recorded at the time of admission. Clinical profile and outcome was assessing with Benzel-Larson Grade included one to seven grades. **Results:** Total 87 patients of cervical spine injury were included in study, 25 patients were included in SCIWORA group while 62 patients were having associated with bony injuries. Both SCIWORA and SCI with bony injury were more commonly due to fall from height and more common in males. Neurological recovery was more in SCIWORA than bony injury patients. **Conclusion:** Cervical spine cord injury is commonly seen in children, while associated bony injury is common in adult population. Early diagnosis and proper management for SCIWORA can improve neurological outcome in selected group of patients.

Keywords: Cervical Spine, SCIWORA (Spinal Cord Injury without Bony Injury).

Corresponding Author: Dr. Vinod Sharma, Associate Professor, Department of Neurosurgery, SMS Medical College, Jaipur, Rajasthan, India.

Received: February 2019

Accepted: February 2019

Introduction

SCIWORA (Spinal cord injury without radiological abnormality) was defined by Pangl & Wilberger in 1982 as the "objective signs of myelopathy as a result of trauma" in which no evidence of fracture, subluxation or instability on plain radiographs or computed tomography (CT) is demonstrated. The SCIWORA is documented in children^{2,3}, probably due to increased elasticity of the paediatric spine. It is not described very well in adults with normal spinal canals. The mechanism for such injuries in adults was initially postulated to be either hyperflexion dislocation with immediate reduction by muscular action or temporary prolapse of the cervical disc after forcible flexion. However, Taylor and Blackwood (1947) and other authors have convincingly demonstrated that the most likely injury mechanism in adult is hyperextension superimposed on pre-existing cervical spondylosis. The present study was planned to evaluate the clinical profile and outcome in patients of cervical spine injury with or without bony injury.

Subjects and Methods

The present study was conducted at department of neurosurgery S. M. S. Medical College, Jaipur over a period of 3 year. S.M.S is a tertiary care super specialty treatment Centre and also a referral center of Rajasthan state. Being the largest medical institute in the region of Rajasthan, it

caters the health needs of entire region as well as neighboring states. All patients of cervical spine injury with or without evidence of bony injury on X-ray were included in this study. In addition to X-ray finding, patient age, sex and MRI cervical spine were also recorded at the time of admission. Initial X-ray cervical spine was looked for -1) type of bony injury 2) height of inter-body space. Initial neurological assessments were done with Benzel-Larson Grade⁴ included one to seven grades. Grade 1 represents complete cord transection with no sensory and motor function while grade VII represents normal neurological status. Patients were looked for any respiratory difficulties and deep vein thrombosis daily. Complication were noted and treated accordingly. Patient neurological recovery was noted according to Benzel-Larson Grade at the time of discharge. Thus, withal above considerations we had a cohort of 87 patients with cervical spine injury. The data was compiled, summarized and analyzed using frequency distribution and percentage proportion.

Results

Total 87 patients of cervical spine injury were included in study, 25 patients were included in SCIWORA group while 62 patients were having associated with bony injuries. Highest incidence of SCIWORA was reported in age group of 0-20 years while SCI with bony injury were present in 21-40 years of age [Table 1].

[Table 2] revealed males were more likely to have SCIWORA compared to females. Spinal cord injury usually associated with bony injury in females. Both SCIWORA and SCI with bony injury were more commonly due to fall from a height, though RTA also constitutes a major bulk in patients with bony injuries. SCIWORA was much less common with other modes of injury [Table 3].

Myelopathy was seen in every patient of SCI whether with or without bony injuries, while central cord syndrome was seen in 7 patients with bony injury and 4 patients of SCIWORA [Table 4]. These were 2 cases of Brown-Sequard Syndrome in patients with SCIWORA and 4 cases in patients with a bony injury with SCI. [Table 4]. On MR, the signal of cord correlated with the type of cord injury. While cord edema was seen in 12 patients, cord contusion was either seen in association with cord edema (2 patients) or with associated disk herniation (8 patients). 3 patients had normal MR and were suspected to have cord concussion [Table 5].

22 patients presented with grade I in the group SCI with bony injury while 5 patients were in grade I at the time of admission in the group SCIWORA [Table 6].

Table 1: Assessment of age in study groups

Age	SCIWORA No. Of Patients	SCI With Bony Injury No. Of Patients
0-20	13(52%)	7(11%)
21-40	9(36%)	25(40%)
41-60	3(12%)	22(36%)
>60	0(0%)	8(13%)

Table 2: Assessment of sex in cervical SCI

Type Of Injury	Male	Female
SCIWORA	24	1
SCI With Bony Injury	45	17

Table 3: Assessment of Mode of Injury

Mode Of Injury	SCIWORA	SCI With Bony Injury
RTI	6	22
FFH	13	26
Penetrating Injury	2	2
Hit By Animal	2	8
Assults	2	4

Table 4: pattern of the neurological deficit at presentation

Deficit	SCIWORA	SC With Bony Injury
Myelopathy	25	62
Central Cord Syndrome	4	7
Anterior Cord Syndrome	1	1
Brown-Sequard Syndrome	2	4
Horners Syndrome	0	0

Table 5: MRI Apperance Of Injured Spinal Cord (Sciwora)

MRI Finding	No. Of Patients	%
Cord Edema	12	48%
Cord Contusion	0	0%
Cord Edema With Contusion	2	8%
Disc Herniation With Or Without Edema And Contusion	8	32%
Normal	3	12%

Table 6: Neurological Statuses At The Time Of Admission And Discharge

Benzel & Larsons grade	At admission		At discharge	
	Sciwora	SCI with bony injury	Sciwora	SCI with bony injury
Grade 1	5	22	2	9
Grade 2	12	21	1	8
Grade 3	7	16	4	10
Grade 4	1	2	2	3
Grade 5	0	1	7	11
Grade 6	0	0	6	7
Grade 7	0	0	1	1
expire	0	0	2	13

Discussion

Our study population was implicitly closed to the age and sex distribution of cervical spine injury in general population. The age range was 0-70 years with a mean of 31.1 years and sex distribution was roughly 4:1, similar to the general population with cervical spine injury as has been reported in earlier studies as well. Approximately 25.7% of the patients have spinal cord injury without bony injury in the present study. The highest incidence of SCIWORA was reported in the age group of 0-20 years. This finding was similar to other studies, as SCIWORA is a disease of the pediatric spine. The common pathogenesis is the inherent elasticity of the soft tissues, which ensures immediate spontaneous reduction after considerable intersegmental displacement. SCIWORA in adults is not well documented. Chen et al.^[5] described five cases with traumatic central cord syndrome and an abnormal cord signal on MRI but without definitive compression of the cord. Pathogenesis for SCIWORA in adult group affirms that degenerative changes or calcification of the posterior common vertebral ligament produce excessive traction of the spinal cord during the accident and can influence the development of SCIWORA,^[6] while others deny this relationship, and believe these changes have no direct influence on the presence of neurological impairment following low-energy trauma.^[7] MRI cervical spine is a useful tool to evaluate SCIWORA and the investigation of choice.^[8] Our study revealed cord edema was seen in 12 patients, cord contusion was either seen in association with cord edema (2 patients) or with associated disk herniation (8 patients). 3 patients had normal MR and were suspected to have cord concussion but when this is not conclusive; diffusion MRI or proton emission tomography (PET) may be useful in those patients with negative MRI and with myelopathy neurological syndrome. Another option to be considered is the use of somatosensory evoked potentials.⁹ Our study revealed maximum patients with SCIWORA presented at Benzel-Larson Grade II as compared to those with a bony injury who mostly presented in Benzel-Larson Grade I. In

SCIWORA patients, at the time of discharge, mostly the patients had improved to a higher grade at Benzel and Larson grade V and Grade VI. 7 patients were discharged at grade V, 6 patients in grade VI while 2 patients expired in the present study. While in patients with bony injuries usually the Benzel and Larson grade at the time of discharge was poor.^[9] patients did not improve and were discharged in grade I. 8 patient in grade II, 10 patients grade III, 13 patients died in SCI with a bony injury in the present study. Above finding indicates spinal cord injury without bony injury had good prognosis compared to spinal cord injury with a bony injury. Although there are several factors that influence the likelihood of neurological recovery in these patients,^[10] and these are primarily related to the initial injury the diameter of the canal the patient's age, the extent of the injury,^[11,12] the presence of disc-ligament lesions and the severity of the neurological syndrome.

Conclusion

SCIWORA is a syndrome that defines posttraumatic SCI in patients with neurological findings without any evidence of fractures or mal-alignment in plain X-rays and CT. cervical spine cord injury is commonly seen in children, while the associated bony injury is common in the adult population. With its superior ability to reveal soft tissue pathologies and prognostic value, MRI is accepted as the imaging modality early diagnosis and proper management for SCIWORA can improve neurological outcome in a selected group of patients.

References

1. Pang D, Wilberger JE. Spinal cord injury without radiographic abnormalities in children. *J Neurosurg* 1982;57:114-29.
2. Pang D, Pollack IF. Spinal cord injury without radiographic abnormality in children: the SCIWORA syndrome. *J Trauma* 1989;29:654-64.
3. Dickman CA, Zabramski JM, Hadley MN, Rekate HL, Sonntag VK. Paediatric spinal cord injury without radiographic abnormalities: report of 26 cases and review of the literature. *J Spinal Disord* 1991;4:296-305.
4. Benzel EC, Larson SJ: Functional recovery after decompressive operation for thoracic and lumbar spine fractures. *Neurosurgery* 19: 772-778, 1986
5. Chen TY, Lee ST, Lui TN, et al. Efficacy of surgical treatment in traumatic central cord syndrome. *Surg Neurol* 1997;48:435-40.
6. Wenger M, Adam PJ, Alarcón F, Markwalder TM. Traumatic cervical instability associated with cord oedema and temporary quadriplegia. *Spinal Cord*. 2003;41(9):521-6.
7. Onishi E, Sakamoto A, Murata S, Matsushita M. Risk factors for acute cervical spinal cord injury associated with ossification of the posterior longitudinal ligament. *Spine (Phila Pa 1976)*. 2012;37(8):660-6.
8. D.Lammertse, D.Dungan, J.Dreisbachetal., "Neuroimaging in traumatic spinal cord injury: An evidence-based review for clinical practice and research," *e Journal of Spinal Cord Medicine*, vol. 30, no. 3, pp. 205–214, 2007.
9. Kasimatis GB, Panagiotopoulos E, Megas P, Matzaroglou C, Gliatis J, Tyllianakis M, et al. The adult spinal cord injury without radiographic abnormalities syndrome: magnetic resonance imaging and clinical findings in adults with spinal cord injuries having normal radiographs and computed tomography studies. *J Trauma*. 2008;65(1):86-93.
10. Yamazaki T, Yanaka K, Fujita K, Kamezaki T, Uemura K, Nose T. Traumatic central cord syndrome: analysis of factors affecting the outcome. *Surg Neurol*. 2005;63(2):95-9.
11. Dolan EJ, Tator CH, Endrenyi L. The value of decompression for acute experimental spinal cord compression injury. *J Neurosurg*. 1980;53(6):749-55.
12. Pavlov H, Torg JS, Robie B, Jahre C. Cervical spinal stenosis: determination with vertebral body ratio method. *Radiology*. 1987;164(3):771-5.

Copyright: © the author(s), 2019. It is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits authors to retain ownership of the copyright for their content, and allow anyone to download, reuse, reprint, modify, distribute and/or copy the content as long as the original authors and source are cited.

How to cite this article: Acharya S, Sharma V, Sharma S, Acharya J. Evaluation of Clinical Profile and Effect of Cervical Spinal Cord Injuries with or Without Bony Injuries. *Asian J. Med. Res.* 2019;8(1):SG08-SG10.
DOI: [dx.doi.org/10.21276/ajmr.2019.8.1.SG3](https://doi.org/10.21276/ajmr.2019.8.1.SG3)

Source of Support: Nil, **Conflict of Interest:** None declared.