

Prevalence of Primary Open Angle Glaucoma in Patients with Myopia.

Archana Pushpakaran¹, Meenu C Jacob², Sedhu P.A³, Cyriac Kurian⁴

¹Associate Professor, Department of Ophthalmology, Sree Narayana institute of Medical Sciences (SNIMS), ²Assistant Professor, Department of Ophthalmology, Sree Narayana institute of Medical Sciences (SNIMS), ³Professor, Department of Ophthalmology, Sree Narayana institute of Medical Sciences (SNIMS), ⁴Professor & HOD, Department of Ophthalmology, Sree Narayana institute of Medical Sciences (SNIMS).

Abstract

Background: Myopia is a condition of the eye that makes it difficult to see distant objects. The present study was conducted to assess prevalence of primary open angle glaucoma in patients with myopia. **Subjects and Methods:** This cross-sectional study was conducted in outpatient Department of Ophthalmology, Sree Narayana institute of Medical Sciences. A minimum of 100 patients with myopia were included in the study. Prevalence of primary open angle glaucoma in patients with myopia was recorded. **Results:** In right eye, patients with IOP between 8-21 is 84%, > 21 is 16%. In left eye, patients with IOP between 8-21 is 85%, > 21 is 15%. In myopic patients, 15% of patients had POAG, 1% with ACG. **Conclusion:** Glaucoma, one of the leading causes of irreversible blindness in the adult population worldwide, is a progressive optic neuropathy. Primary open angle glaucoma (POAG) is the most commonly reported type of glaucoma in population based prevalence studies worldwide. Elevated intraocular pressure is a well-known major risk factor for POAG.

Keywords: Glaucoma, intraocular pressure, Primary open angle glaucoma.

Corresponding Author: Dr. Meenu C Jacob, Assistant Professor, Department of Ophthalmology, Sree Narayana institute of Medical Sciences (SNIMS).

Received: May 2019

Accepted: May 2019

Introduction

Myopia is a condition of the eye that makes it difficult to see objects that are far away without optical correction. Although myopia can be induced by lens or corneal curvature, or by other factors, axially elongated eyes represent a major portion of all myopia cases. Axial elongation can affect the intraocular structure, such as the optic disc or macula, which are target sites for glaucomatous damage. Many studies have reported that myopia is a risk factor for glaucoma development. Large-scale population based studies have also reported that myopia is a risk factor for glaucoma development.^[1,2]

Primary open angle glaucoma (POAG) is a chronic progressive optic neuropathy. Direct and convincing evidences for primary mechanisms of glaucoma are still lacking and early detection or predicting progression of POAG remains difficult and challenging.^[3]

Many studies have investigated and reported risk factors associated with glaucoma. Elevated intraocular pressure (IOP) is a well-known major risk factor for POAG. Evidence shows that lowering IOP reduces the risk of development or slows the progression of glaucoma. In addition, there is growing evidence that other risk factors like age, gender, race, refractive errors, heredity and systemic factors may play a role in glaucoma pathogenesis.^[4]

Many studies found that high myopia has been associated with POAG. It is possible that myopic individuals may be at

increased risk for the development of glaucoma. Epidemiologic evidence suggests that high myopia is a risk factor for the development and the progression of glaucomatous optic neuropathy.^[5]

Subjects and Methods

This cross-sectional study was conducted in outpatient Department of Ophthalmology, Sree Narayana institute of Medical Sciences. A minimum of 100 patients with myopia were included in the study. The study was approved by the institutional ethics review board. All patients diagnosed with myopic refractive error > 3.00 diopter. A detailed physical examination and ophthalmological examination was done.

During Ophthalmological examination, best corrected visual acuity (BCVA) was assessed using an illuminattx1 Snellen's chart, with the patient seated at 6 meters distance. Near vision was assessed, using Jaeger's near vision chart. Colour vision was checked using Ishihara's pseudo-isochromatic charts. BCVA was checked by Skiascopy or streak retinoscopy. Slit lamp examination was performed to rule out anterior segment pathology. Gonioscopy was done with Goldmann's three mirror gonioscope and the anterior chamber angle was graded according to modified Shaffer's grading.

Dilated fundus examination by indirect ophthalmoscopy, followed by a slit lamp biomicroscopic evaluation with 78 D lens and 90 D lens to evaluate the posterior pole

including the optic disc was done. Visual field was tested using the Automated Humphrey Visual Field Analyzer. The test algorithm used was SITA standard 30-2. The visual fields were analyzed as per the Anderrson Patella criteria and the severity of the glaucomatous field changes were graded according to the Hodapp, Parrish and Andersson classification guidelines given in the European Glaucoma Society guidelines for Glaucoma, 3a Edition. Patients are analyzed for some parameters such as C:D ratio, IOP, visual fields, each eye was analyzed separately as there could be differences between the two eyes of the same patient which would be missed if both eyes were evaluated together. Data are presented as mean and standard deviation for continuous variables. Statistical significance was considered when P was < 0.05.

Results

Table 1: Distribution of patients

Total- 100		
Gender	Males	Females
Number	51	49

[Table 1] shows that out of 100, males were 51 and females were 49.

Table 2: Best corrected visual acuity in RE.

BCVA	MYOPIA	
	Count	%
PL +ve	0	0.0
HM+	1	1.0
CF 1m	2	2.0
CF 2m	1	1.0
CF 3m	1	1.0
3/60	2	2.0
6/60	4	4.0
6/36	2	2.0
6/24	2	2.0
6/12	3	3.0
6/9	25	25.0
6/6	57	57.0
Total	100	100.0

[Table 2] shows best corrected visual acuity in RE & LE.

Table 3: Best corrected visual acuity LE

BCVA LE	MYOPIA	
	Count	%
HM+	2	2.0
CF 3m	2	2.0
1/60	1	1.0
3/60	2	2.0
6/60	3	3.0
6/36	3	3.0
6/24	1	1.0
6/18	1	1.0
6/12	0	0.0
6/9	22	22.0
6/6	63	63.0
Total	100	100.0

[Table 3] shows best corrected visual acuity LE.

Table 4: IOP variation in RE & LE




Figure 1:

[Figure 1] shows that in right eye, patients with IOP between 8-21 is 84%, > 21 is 16%. In left eye, patients with IOP between 8-21 is 85%, > 21 is 15%.

Table 5: Prevalence of POAG

Classification	MYOPIA	
	Count	%
	84	84.0
ACG	1	1.0
POAG	15	15.0
Total	100	100.0

[Table 5] shows that in myopic patients, 15% of patients had POAG, 1% with ACG.

Discussion

The World Health Organization has estimated that 8.9 million people in India are blind, of which 12.8% are due to glaucoma. The visual disability and irreversible blindness from glaucoma has significant socio- economic impact and the problem is expected to reach alarming proportions in few years. Risk for blindness from Primary Open Angle Glaucoma is high because of advanced stage at the time of diagnosis, onset of glaucoma at young age, inadequate intraocular pressure control, high rate of progression despite treatment, undiagnosed glaucoma and missed opportunities for diagnosing glaucoma.^[6]

Screening for glaucoma in the population will help to identify cases early. However it may be economically unviable to screen the entire population. Identification of the risk factors will help to select high risk patients for screening which in turn should lead to an overall reduction in the blindness due to this disease and the morbidity associated with it.^[7]

Available data suggests that the prevalence of POAG varies from race to race and is influenced by various factors like age, gender, and other associated risk factors. The prevalence of POAG among adult black populations is much higher than the prevalence in predominately white adult populations which ranges from 1.1 to 3%. The prevalence estimated for POAG, in east Asia varies from 0.5 to 2.3% and from India is between 0.41% to 2.56%.^[8]

Our study is a hospital based study. It was performed to calculate the prevalence of POAG in patients with myopia, 100 myopic patients were included. Population based studies done in rural and urban south India 82'83 and in Central India also did not show any significant gender variation between patients with or without POAG. The blue mountain eye disease study showed a higher incidence of POAG in women than in men¹³, where the odds ratio of females having POAG was 1.3 (95% CI, 0.7, 2.6), although this was not statistically significant. The study done by Faschinger et al,^[8] in Barbados which showed that males were more likely to have POAG.

In our study 15% of myopic (> 3D to 6D) population between the age group 20-40 years developed POAG (p<0.05). The Blue Mountain study⁵⁷ which is a population based cross sectional study showed glaucoma was present in 4.2% of eyes with low myopia (> or = -1.00 D to < -3.0 D) and 4.4% of eyes with moderate to high myopia (> or = -3.0D) as compared to 1.5% of eyes without myopia. This study concluded that Myopic subjects had a twofold to three fold increased risk of glaucoma compared with that of non myopic subjects. The risk was independent of other glaucoma risk factors and IOP. Our results were in similar to Aravind study,^[9] which showed myopia had a statistically significant association with POAG. Myopic subjects had a twofold or threefold increased risk of glaucoma compared with that of non myopic subjects.

Population based study by Chang RT,^[10] did not show any association between myopia and POAG. The Limitation of this study is that it is a hospital based cross-sectional study as compared to the other related studies which have been done which were mainly population based epidemiological studies. Another drawback is the small sample size as compared to many of the other studies.

Conclusion

Glaucoma, one of the leading causes of irreversible blindness in the adult population worldwide, is a progressive optic neuropathy. Primary open angle glaucoma (POAG) is the most commonly reported type of glaucoma in population based prevalence studies worldwide. Elevated intraocular pressure is a well-known major risk factor for POAG.

References

1. Loyo-Berrios NI, Blustein JN. Primary open glaucoma and myopia: A narrative review. 2007;106(2):85-91-95.
2. Kim NR, Lee ES, Seong GJ, Kang SY, Kim JH, Hong S, Kim CY. Comparing the ganglion cell complex and retinal nerve fibre layer measurements by Fourier domain OCT to detect glaucoma in high myopia. J Glaucoma 2011;95(8):1115-1121
3. Cedrone C, Mancino R, Cerulli A, Cesareo M, Nucci C. Epidemiology of primary glaucoma: prevalence, incidence, and blinding effects. J Glaucoma 2008;173:3-14.
4. Marcus MW, de Vries MM, Junoy Montolio FG, Jansoni NM. Myopia as a risk factor for open angle glaucoma: A systematic review and meta-analysis. J Glaucoma 2011;118(10):1989-94 e2
5. Sia DI, Edussuriya K, Sennanayake S, Senaratne T, Selva D, Casson RJ. Prevalence of and risk factors for primary open angle glaucoma in central Sri Lanka: the Kandy eye study. J Glaucoma. 2010;17 (4): 211-216.
6. McMonnies CW. Intraocular pressure spikes in keratectasia, axial myopia, and glaucoma. Invest Ophthalmol Vis Sci. 2008;85(10):1018-1026
7. Blumen Ohana E, Blumen MB, Bluwol E, Derri M, Chabolle F, Nordmann JP. Primary open angle glaucoma and snoring: prevalence of OSAS. 2010;127(5):159-164.
8. Faschinger C, Mossbock G. Myopia and glaucoma. Invest Ophthalmol Vis Sci. 2007;157(7-8):173-177.
9. Aravind, Suzuki Y, Iwase A, Araie M. Risk factors for open-angle glaucoma in a Japanese population: the Tajimi Study. Ophthalmology. 2006; 113: 1613-1617.
10. Chang RT. Myopia and glaucoma. Invest Ophthalmol Vis Sci. 2011;51 (3): 53-63

Copyright: © the author(s), 2019. It is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits authors to retain ownership of the copyright for their content, and allow anyone to download, reuse, reprint, modify, distribute and/or copy the content as long as the original authors and source are cited.

How to cite this article: Pushpakaran A, Jacob MC, Sedhu PA, Kurian C. Prevalence of Primary Open Angle Glaucoma in Patients with Myopia. Asian J. Med. Res. 2019;8(2):OT01-OT03.
DOI: dx.doi.org/10.21276/ajmr.2019.8.2.OT1

Source of Support: Nil, **Conflict of Interest:** None declared.